



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

*Foreign statistical reports of countries and cities—Yearly and monthly.*

AFRICA—*Lorenzo Marquez*.—Month of March, 1900. Estimated population, 4,000. Total number of deaths, 69, including 1 from tuberculosis.

AUSTRALIA—*New South Wales—Sydney*.—Month of June, 1900. Estimated population, 70,000. Total number of deaths, 47, including diphtheria, 11; enteric fever, 4; measles, 2; scarlet fever, 9, and 2 from typhus.

BRITISH HONDURAS—*Belize*.—Month of June, 1900. Estimated population, 30,000. Total number of deaths, 47, including enteric fever, 1; whooping cough, 5, and 4 from phthisis pulmonalis.

GERMANY—*Stuttgart*.—Month of February, 1900. Estimated population, 179,328. Total number of deaths, 269, including diphtheria, 6; enteric fever, 1; scarlet fever, 2, and 27 from phthisis pulmonalis.

ITALY—*Florence*.—Month of March, 1900. Estimated population, 201,425. Total number of deaths, 371, including diphtheria, 4; enteric fever, 1; measles, 2; scarlet fever, 1, and 59 from tuberculosis.

*Genoa*.—Month of May, 1900. Estimated population, 226,432. Total number of deaths, 289, including diphtheria, 2; enteric fever, 3; measles, 6; whooping cough, 1; and 43 from tuberculosis.

NORFOLK ISLAND.—Months of May and June, 1900. Estimated population, 846. Total number of deaths, 2. No deaths from contagious diseases.

SWITZERLAND.—Reports for the two weeks ended August 4, 1900, from 15 cities and towns having an aggregate estimated population of 691,000, show a total of 510 deaths, including diphtheria, 5; enteric fever, 7; measles, 6; whooping cough, 7; and 54 from phthisis pulmonalis.